



PATIENT INFORMATION

Name, Parent Name, Date of Birth, Sex, Address, City, State, Zip, Telephone, Date of Specimen

BILLING INFORMATION

Insurance, Patient, Lab, Doctor, FOR INSURANCE BILLING, PROVIDE COMPLETE INSURANCE INFORMATION AND SEND PHOTOCOPY OF PATIENT'S INSURANCE CARD, Inpatient, Outpatient, P.O. #, Facility, Address, City, State, Zip, Date of Admission, MR/Lab #, Insurance Company, Policy #, Group #, Authorization #

REQUESTING DOCTOR

Last Name, First Name, Practice/Facility Name, Address, City, State, Zip, Telephone, HIPAA Fax, Email, NPI #

Physician Signature Required by CMS

PLEASE COMPLETE ALL ITEMS – TYPE / PRINT PER HIPAA REGULATIONS

KSL Beutner Laboratories complies with the Health Insurance Portability and Accountability Act (HIPAA). The information provided herein will remain strictly confidential.

COMPLETE BILLING AUTHORIZATION ON BACK BEFORE SUBMITTING

REQUEST FREE TEST KITS AT 1-800-288-0549, INDICATE BELOW, OR ORDER AT BEUTNERLABS.COM


BIOPSY KITS, SERUM KITS, 4-TUBE KITS, #

DATA ON SKIN AND MUCOSAL BIOPSIES – FOR ALL DIRECT IF SPECIMENS, PLEASE FILL OUT THIS PANEL

Interpretation of direct immunofluorescence (IF) of tissue specimens requires knowledge of the location of the biopsy relative to the clinical lesion.

Biopsy of: Specific location, Lesional site, Normal site, Perilesional site

For Lupus: sun exposed, sun protected, Please draw the biopsy site as it relates to the lesion.



Anti Ro(SS-A)/La(SS-B), ANA titer/pattern

Fixative used: Beutner Labs, Other

Clinical diagnoses

ICD-10 Code (Required)

Clinical findings

SUGGESTED BIOPSY SITES FOR DIRECT IMMUNOFLUORESCENCE AND SERUM STUDIES

BULLOUS DISEASES: INDICATED DIRECT IMMUNOFLUORESCENT (DIF) AND SERUM STUDIES SKIN LESIONS

If pemphigoid or epidermolysis bullosa acquisita is suspected, take skin biopsy with ~2/3 normal skin and ~1/3 edge of lesion. If pemphigus is suspected, take skin biopsy with ~2/3 normal skin and ~1/3 lesion edge plus serum for best diagnostic results. If dermatitis herpetiformis is suspected, take normal skin ~3 mm from lesion for best results. If porphyria or pseudoporphyria is suspected, take skin biopsy with ~2/3 normal skin and ~1/3 lesion edge for best DIF results. If in doubt, take two biopsies – one perilesional, as for pemphigoid, and one normal, as for DH, for best results. If eruptions with other non-disease specific immune deposits are suspected, including lichenoid planus or lichenoid eruption or related disorders, take biopsy as for porphyria for DIF and lesional biopsy for light microscopy.\*

MUCOSAL LESIONS

If pemphigoid is suspected, take normal mucosa ~3 mm from lesion or Nikolsky sign. If pemphigus or paraneoplastic pemphigus is suspected, take normal mucosa ~3 mm from lesion or Nikolsky sign plus serum. If erosive lichen planus (LP) is suspected, take mucosal biopsy with ~2/3 normal mucosa and ~1/3 edge of lesion or of Nikolsky sign for best DIF results. HEREDITARY EPIDERMOLYSIS BULLOSA (EB) If hereditary EB needs to be classified or confirmed, take biopsy of induced lesion in normal skin.

CONNECTIVE TISSUE DISEASES: INDICATED DIRECT IMMUNOFLUORESCENT (DIF) AND SERUM STUDIES

If SLE is suspected, take biopsy of sun-exposed normal skin of forearm for DIF for LE band test and blood in serum separator tube (SST). Serum can be sent in red top tube. If DLE is suspected, take biopsy of a sun-exposed lesion of 3 or more months duration for DIF and for light microscopy.\* Non-sun-exposed areas are of little value.

If SCLE or Sjögren Syndrome is suspected, take sun-exposed skin lesion biopsy for DIF for in vivo ANA and blood in SST for serum tests. Serum can be sent in red top tube.

If systemic sclerosis (SSc) is suspected, take biopsy of sun-exposed skin for DIF studies and blood in SST for serum tests. Serum can be sent in a red top tube in red top tube

IMMUNE COMPLEX MEDIATED VASCULITIS: INDICATED DIRECT IMMUNOFLUORESCENT STUDIES\*

If leukocytoclastic vasculitis is suspected, (or most other immune complex vasculitides), take biopsy for DIF of a fresh lesion, less than 48 hours old. A lesional biopsy for light microscopy is also indicated. If Henoch Schoenlein purpura is suspected, take biopsy of a fresh lesion (less than 48 hours old).

\*Additional/separate biopsy should be sent in formalin for light microscopy studies. Light microscopic studies of lesional biopsy sites are indicated, in most cases of vesiculobullous, connective tissue disease, and vascular disorders.


## BULLOUS DISEASES

### SKIN AND MUCOSAL BIOPSY STUDIES

- 001  Direct immunofluorescence (DIF) for IgG, IgA, IgM, fibrin and C3. If needed, IgG4 and/or IgG1 are also added for greater sensitivity.
- 002  Differentiation of bullous pemphigoid from epidermolysis bullosa acquisita by DIF of 1 M NaCl split biopsy (with no vesicles)
- 003  Light microscopy (H&E)

#### BLISTERING AND OTHER ERUPTIONS

Please fill in data on front of form. See front of form for abbreviations.



- (1) Skin biopsy in most pemphigus/pemphigoid cases.  
(2) Mucosal or skin biopsy for DH; 2nd skin biopsy for pemphigus/pemphigoid cases.

### SERUM STUDIES

- 010  Laminin 332 (Laminin 5 or epiligrin) antibodies (IIF)
- 014  Split skin, pemphigoid/epidermolysis bullosa acquisita (EBA) differentiation by IgG & IgG4 indirect IF (Split-IIF)
- 015  Desmoglein (Dsg3 and Dsg1) ELISA
- 016  BP230 (BPAG1) and BP180 (BPAG2) ELISA
- 017  Paraneoplastic pemphigus antibody titer (on rat bladder) (PPA-IIF)
- 009  Envoplakin antibodies (ELISA)
- 018  Pemphigus antibody titer prognostic test comparing old and new sera for IgG and IgG4 (Monkey esophagus-IIF)
- 023  Collagen VII antibodies (ELISA)
- 024  IgA split skin - IIF, linear IgA bullous dermatosis (LABD)

### SERUM PANELS

#### BASIC PEMPHIGUS - PEMPHIGOID SCREEN

- 013  Monkey esophagus for IgG and IgG4 antibodies by indirect immunofluorescence (IIF)

#### PEMPHIGUS

- 025  Monkey esophagus for IgG and IgG4 antibodies by IIF (013) and Dsg 3/1 ELISA (015)

#### PARANEOPLASTIC PEMPHIGUS

- 026  Monkey esophagus for IgG, IgG1 & IgG4 antibodies by IIF (013) plus IIF on rat bladder (017) and Dsg3/1 ELISA (015), envoplakin ELISA (009), and A2ML1 Immunoblot\*

**SUB EPIDERMAL BULLOUS DISEASES – All panels below include IIF tests on monkey esophagus (#013) and Split Skin (#014). KSL Beutner Laboratories may perform relevant reflex tests based on results of IIF.**

#### PEMPHIGOID

- 027\*  Monkey esophagus (013) and split skin for IgG & IgG4 antibodies by IIF (014) and BP180 plus BP230 ELISA (016)

#### EBA/BULLOUS LE

- 061\*  Monkey Esophagus (013), Split Skin for IgG and IgG4 by IIF (014) and Collagen VII antibodies by ELISA (023)

#### LAMININ 332 PEMPHIGOID

- 062\*  Monkey Esophagus (013), Split Skin for IgG and IgG4 (014), and Laminin 332 (Laminin 5 or epiligrin) antibodies by IIF (010)

#### DERMATITIS HERPETIFORMIS & CELIAC DISEASE

- 020  IgA/IgG anti endomysial antibodies (AEmA-IIF)
- 022  IgA epidermal transglutaminase antibodies (eTG-ELISA)
- 053  IgA/IgG Tissue Transglutaminase (tTG-ELISA)
- 054  IgA/IgG Gliadin Peptides (ELISA)
- 055  Comprehensive Dermatitis Herpetiformis/Celiac Disease Panel – IgA/IgG Anti-Endomysial Antibodies (AEmA-IIF) (020), IgA Epidermal Transglutaminase Antibodies (eTG-ELISA) (022), IgA/IgG Tissue Transglutaminase (tTG-ELISA) (053), IgA/IgG Gliadin Peptides (ELISA) (054)

\*Turnaround time may be longer for this test.

## BILLING AUTHORIZATION

KSL Beutner Laboratories will bill your medical insurance using the information provided to us on this Test Request Form. If we do not receive all the required insurance information, you will receive a bill directly from KSL Beutner Laboratories. Please be aware that we may not be participating with your insurance plan and that insurance payment may vary based on your coverage. **See "Billing and Insurance" page in the "Resources" drop-down menu at [www.beutnerlabs.com](http://www.beutnerlabs.com) for a list of participating insurances or to request self-pay pricing.** Ultimately, you are responsible for the full payment or balances not covered by your insurance.

**I certify that I have read and understand the information above and consent to the testing procedure(s). I understand that my test(s) are being sent to KSL Beutner Laboratories for analysis and I accept full financial responsibility for any payment that may not be covered by my insurance. I authorize KSL Beutner Laboratories to release medical reports to my health insurance as necessary to process insurance claims and I authorize my insurance to pay KSL Beutner Laboratories directly.**

SIGNATURE OF PATIENT, LEGAL GUARDIAN, OR POWER OF ATTORNEY

X

SIGNATURE ON FILE

DATE

## BULLOUS DISEASES

### HEREDITARY EPIDERMOLYSIS BULLOSA

- 011  IF mapping of pencil eraser induced clefts in normal skin away from lesions for hereditary epidermolysis bullosa (EB): (a) Primary screen for collagen type VII, collagen type IV, & keratin 14 to check for suitable biopsy; (b) For suitable biopsies, test for collagen XVII, alpha6 integrin, beta4 integrin, laminin 332 and plectin to differentiate dystrophic EB and junctional EB from EB simplex

## LUPUS ERYTHEMATOSUS, CONNECTIVE TISSUE & VASCULAR DISEASE


### SKIN AND MUCOSAL BIOPSIES

- 005  Direct immunofluorescence for SLE, DLE & SCLE with tests for IgG, IgA, IgM, fibrin and C3
- 007  Direct immunofluorescence for dermatomyositis with tests for IgG, IgA, IgM, fibrin, C3 and C5b-9
- 006  Direct immunofluorescence for vasculitis with tests for IgG, IgA, IgM, fibrin and C3
- 003  Light microscopy (H&E) to rule out LE

#### COLLAGEN VASCULAR DISEASES

Please fill in data on front of form. See front of form for abbreviations.

- (3) Sun exposed skin biopsy in most LE cases. Skin biopsy to rule out Henoch Schoenlein purpura and vasculitis (lesion < 48 hours old).  
(4) Sun-exposed skin biopsy to rule out SLE.



### CHRONIC ULCERATIVE STOMATITIS

- 033  Stratified epithelium specific ANA (SES-ANA), Monkey esophagus, and ANA (HEp-2)

## SYSTEMIC CONNECTIVE TISSUE DISEASE SERUM STUDY PANELS

### ANA SCREEN

- 042  ANA titer and pattern on HEp-2 cells (IIF)

### SLE PANEL

- 043  ANA titer and pattern on HEp-2 cells (IIF), antibodies to dsDNA, Ro (SS-A), La (SS-B), Sm, RNP, Ribo P (ELISA)

### MIXED CONNECTIVE DISEASE PANEL

- 044  ANA titer and pattern on HEp-2 cells (IIF), Sm, RNP, Jo-1 and Scl-70 (ELISA)

### SJÖGREN SYNDROME PANEL

- 045  ANA titer and pattern on HEp-2 cells (IIF), Rheumatoid Factor, Ro (SS-A), La (SS-B), CENP (ELISA)

### DRUG INDUCED LUPUS PANEL

- 046  ANA titer and pattern on HEp-2 cells (IIF), antibodies to dsDNA, Ro (SS-A), La (SS-B), Sm, Histone (ELISA)

### SYSTEMIC SCLEROSIS PANEL

- 047  ANA titer and pattern on HEp-2 cells (IIF), RNP, Scl-70, CENP, RNA Polymerase III (ELISA)

## ADDITIONAL DISEASE PANELS

### RHEUMATOID ARTHRITIS

- 056  Cyclic citrullinated peptide (CCP), Rheumatoid Factor (RF), SR-A (ELISA)

### VASCULITIS

- 057  Anti-Neutrophil Cytoplasmic Antibody (ANCA- IIF), Myeloperoxidase Ab (MPO), Proteinase 3 Ab (PR3) (ELISA)

**NOTE: Serum studies / profiles require a minimum of 2-5ml of serum**